

Table 1—Sample of SRS data table (dummy data)

Species	Life zone	Area <i>acres</i>	Height		
			Actual	Median	Mean
Hardwood					
Big-leaf mahogany	Tropical dry and moist	.20	00.1 – 112.5	7 million fibers	Trace – .32
	Tropical moist	31.12	1,000,000 – 0.01	None	18.4 – 550
	Premontane rain	—	155	1.3	0.1 > 38.5
	Tropical dry	<u>100.05</u>	> 60	.004	Average daily
Total		131.37			
Total species		131.37			

— = Not applicable.

Table 2—Sample of SRS data table with metric units (dummy data)

Forest type	Land area		Range	Location
	<i>km</i> ²	%	<i>hours</i>	
Orey ^a	450	1.3	1,050	Streams and rivers in Wyoming and Montana
Orey + other species	170	0.5	300	
Cativo ^b	300	0.8	—	
Cativo + other species	195	0.5	0.03	Newfoundland, Canada
Mixed forests	17 880	50.4	—	
Mangroves	1 760	5	100	NG
Protection forests	14 740	41.5	3	NR
Total	35 497	100.0		

— = Not applicable; NG = not given; NR = no reports found.

^a *Campnosperma panamensis*.

^b *Priora copaiifera*.

Source: Harcourt and Sayer (1996), INRENARE (1990).